



Faculty of Pharmacy Department of Pharmacognosy & Natural Products Chemistry Panepistimiopolis Zografou 15771, Athens

Tel: +30 210 72 74052 magiatis@pharm.uoa.gr



Athens, 21/10/2020 Cert.Num: 2021-C00007

CERTIFICATE OF ANALYSIS

Analysis Date: 21/10/2020

Owner:

ARGALI

Variety:

KORONEIKI

Origin:

GARGALIANOI AGIOS DIMITRIOS MESSINIA GREECE

Chemical Analysis

Oleocanthal	287	mg/Kg
Oleacein	218	mg/Kg
Oleocanthal + Oleacein (index D1)	504	mg/Kg
Ligstroside aglycon (monoaldehyde form)	135	mg/Kg
Oleuropein aglycon (monoaldehyde form)	173	mg/Kg
Ligstroside aglycon (dialdehyde form)	386	mg/Kg
Oleuropein aglycon (dialdehyde form)	195	mg/Kg
Total tyrosol derivatives	807	mg/Kg
Total hydroxytyrosol derivatives	585	mg/Kg
Total polyphenols analyzed	1.392	mg/Kg

Comments:

The levels of oleocanthal and oleacein are higher than the average values (135 and 105 mg/Kg respectively) of the sample included in the international study performed at the University of California, Davis.

The daily consumption of 20 g of the analyzed olive oil provides 27.8 mg of hydroxytyrosol, tyrosol or their derivatives. Olive oils that contain >5 mg per 20 gr belong to the category of oils that protect the blood lipids from oxidative stress according to the Regulation 432/2012 of the European Union.

It should be noted that oleocanthal and oleacein present important biological activity and they have been related with anti-inflammatory, antioxidant, cardioprotective and neuroprotective activity.

The chemical analysis was performed according to the method published in J.Agric. Food Chem., 2012, 60 (47), pp 11696-11703, J.Agric. Food Chem., 2014 62 (3), 600-607 and OLIVAE, 2015, 122, 22-33.

*Oleomissional+Oleuropeindial **Ligstrodial+Oleokoronal

Magiatis Prokopios

PROKOPIOS MAGIATIS

UNIVERSITY OF ATHENS

DEPARTMENT OF PHARMACOGNOSY
AND NATURAL PROPERTY OF CHEMISTRY